

CLINICAL CASE STUDY

PROBIOTICS AND WEIGHT MANAGEMENT


Investigating the Effect of a Probiotic on Body Composition and Weight Management in Overweight Adults

Executive Summary

Visceral adipose tissue accumulation is currently an important research target because of its connection to metabolic disturbances, cardiovascular disease and type 2 diabetes. This study illustrates how Atlantia designed and conducted a study to investigate the outcome of probiotics supplementation in body composition and weight management assessed through DXA scans.

About the Sponsor

The sponsor is a trusted and pioneer manufacturer of Premium Probiotics. Their dedicated in-house research teams work to formulate probiotics that support desired structure-function claims, but they required

A light green graphic of a molecular structure, consisting of several interconnected hexagons and circles, is positioned behind the text. The text is written in a light green, italicized sans-serif font.

The company had conducted a previous trial; however, the study was insufficiently powered to reach statistical significance in several key outcomes.

DO YOU WANT TO KNOW MORE?

Keep reading about the challenges and objectives



CLINICAL CASE STUDY

Investigating the Effect of a Probiotic on Body Composition and Weight Management in Overweight Adults



a partner in order to design and conduct a gold-standard trial. The sponsor company has conducted a previous trial; however, the study was insufficiently powered to reach statistical significance between groups in several key outcomes.


Challenges and Objectives

The rationale for the current study is to perform a follow-up randomized controlled trial to continue to build the evidence for this probiotic strain in the area of weight management. The sponsors needed a trusted partner able to access to the required study population, with enough expertise to design and conduct a gold-standard trial with probiotics to achieve the following objectives:

- To determine the effect of daily administration of a probiotic strain on Visceral Adipose Tissue (VAT), as well as on other secondary biomarkers and efficacy parameters.
- To determine the effect of a daily administration of a probiotic strain on faecal characteristics.

How Atlantia's Solution Helped

Atlantia's world-class team of nutritionists and research nurses worked closely with the Sponsors' researchers to design and deliver the study objectives. Ireland has one of the highest rates of obesity in Europe. The percentage of people who were classified as overweight or obese in 2015 was 60% (Irish Health Survey), and this has risen to 62% in 2017 (CSO). The research clinic that the company owns and operates in Cork seemed a perfect fit to recruit overweight subjects within the sponsors expected timeline. The company's expertise measuring VAT by DXA was a key factor on the conduction of this project.



CLINICAL CASE STUDY

Investigating the Effect of a Probiotic on Body Composition and Weight Management in Overweight Adults



About the Study

This study was a single-centre, randomised, double-blind, placebo-controlled, parallel study in overweight volunteers. The total population was of 126 participants aged between 25 and 65 years old with a body mass index between 25.0 to 29.9 Kg/m², a waist-hip ratio of ≥ 0.91 for males and ≥ 0.81 for females, having a sedentary lifestyle (exercising less than 2 times per week). The participants should be outside the healthy parameters of visceral adipose tissue, defined as 762 cm³ for males and 256 cm³ for females, assessed by DXA scans. Other exclusion criteria were also considered. Participants were recruited through Atlantia's database, social media platforms, General Practitioners offices and advertisements in local media.

Participants who met the eligibility criteria were randomised (ratio 1:1) to one of the two treatment groups prior writing informed consent: a capsule containing the optimal dose as per the Sponsors pre-clinical results and placebo. Each eligible participant was assigned a randomisation number in chronological order, which corresponds to one of the two groups. The relationship between the randomisation number and the group assignment was unknown to the clinical research team, the sponsor, study site staff, or the participants.

The study involved 5 visits over a 13-15-week period, per participant, including a run-in phase and the intervention period. Participants were instructed to consume one capsule per day, before breakfast for the duration of the study, they were retrained if their compliance was not suitable.

A fasting blood sample (21mls) was collected at various visits. Additional plasma and serum were stored at -80oC at Atlantia's facilities, for future exploratory analysis of inflammatory and satiety biomarkers. Participants will be provided with a stool collection kit and instructions for collecting and storing at home. Participants returned the stool sample and they were stored at -80oC, for future exploratory analysis including shotgun sequencing, SCFA analysis and detection of the strain.

Future plans and Return on Investment

This trial data is still under review and details on the results will be given at another time. The sponsor was highly satisfied with Atlantia's management of the trial from design to conduction.

TO ENQUIRE ABOUT STUDIES SUCH AS THIS

Or in the many other health areas in which we work, please contact our expert team:



Shauna Cusack
Sales Executive
+353 879220160
scusack@atlantiafoodtrials.com



Barry Skillington
Chief Commercial Officer
+353 86 8181543
bskillington@atlantiafoodtrials.com

DO YOU WANT TO KNOW MORE?

Keep reading about the study and its results

